

A second-order difference scheme for the time fractional substantial diffusion equation

Zhaopeng Hao^a, Wanrong Cao^{a,*}, Guang Lin^b

^a*Department of Mathematics, Southeast University, Nanjing 210096, P.R. China.*

^b*Department of Mathematics, School of Mechanical Engineering, Purdue University, West Lafayette, IN 47907, USA.*

Abstract

In this work, a second-order approximation of the fractional substantial derivative is presented by considering a modified shifted substantial Grünwald formula and its asymptotic expansion. Moreover, the proposed approximation is applied to a fractional diffusion equation with fractional substantial derivative in time. With the use of the fourth-order compact scheme in space, we give a fully discrete Grünwald-Letnikov-formula-based compact difference scheme and prove its stability and convergence by the energy method under smooth assumptions. In addition, the problem with nonsmooth solution is also discussed, and an improved algorithm is proposed to deal with the singularity of the fractional substantial derivative. Numerical examples show the reliability and efficiency of the scheme.

Keywords: high-order finite difference method, fractional substantial derivative, weighted average operator, stability analysis, nonsmooth solution

MSC subject classifications: 26A33, 65M06, 65M12, 65M55, 65T50

1. Introduction

Anomalous sub-diffusion process, commonly described by the continuous time random walks (CTRWs), and also known as non-Brownian sub-diffusion, arises in numerous physical, chemical and biological systems; see [3, 5, 13, 21]. The Feynman-Kac formula named after Richard Feynman and Mark Kac, establishes a link between parabolic partial differential equations (PDEs) and Brownian functionals. To figure out the probability density function (PDF) of some

*Corresponding author. Tel/Fax: +86 25 52090590

Email addresses: 230139121@seu.edu.cn (Zhaopeng Hao), wrcao@seu.edu.cn (Wanrong Cao), guanglin@purdue.edu (Guang Lin)

non-Brownian functionals, the fractional Feynman-Kac equation has been derived in [8, 9, 17, 33, 38]. The non-Brownian functionals can be defined by $A = \int_0^t U(x(\tau))d\tau$, where $x(t)$ is the trajectory of a non-Brownian particle and different choices of $U(x)$ can depict diverse systems. In particular, if taking $U(x) \equiv 0$, the fractional Feynman-Kac equation reduces to the well-known fractional Fokker-Planck equation; see [8, 9, 28] for details. Lévy walks give a proper dynamical description in the superdiffusive domain, where the temporal and spatial variables of Lévy walks are strongly correlated and the PDFs of waiting time and jump length are spatiotemporal coupling [33]. Thus, an important operator, fractional substantial derivative has been proposed to describe the CTRW models with coupling PDFs. This spatiotemporal coupling operator was also presented in [17], where the CTRW model with position-velocity coupling PDF was discussed. Recently, Carmi and Barkai [8] also used the substantial derivative to derive the forward and backward fractional Feynman-Kac equations. Due to its potential properties and wide application, the fractional substantial derivative has attracted many scholars' attention; see [6, 11] and references therein.

The fractional substantial derivative operator of order α ($n - 1 < \alpha < n$) is defined by [11]

$${}_a D_t^{\alpha, \lambda} f(t) = {}_a D_t^{n, \lambda} [{}_a I_t^{n-\alpha, \lambda} f](t), \quad {}_a I_t^{\alpha, \lambda} f(t) = \frac{1}{\Gamma(\alpha)} \int_a^t \frac{f(\xi)}{(t-\xi)^{1-\alpha} e^{\lambda(t-\xi)}} d\xi; \quad (1.1)$$

where $\alpha > 0$, λ is a constant or a function independent of variable t , ${}_a I_t^{\alpha, \lambda}$ denotes the fractional substantial operator, and

$${}_a D_t^{n, \lambda} = \left(\frac{d}{dt} + \lambda\right)^n = \left(\frac{d}{dt} + \lambda\right)\left(\frac{d}{dt} + \lambda\right) \cdots \left(\frac{d}{dt} + \lambda\right).$$

It is noted that if λ is a nonnegative constant, then the fractional substantial derivative is equivalent to the Riemann-Liouville tempered derivative defined in [2, 4, 10], and taking $\lambda = 0$ in (1.1) leads to the left Riemann-Liouville derivative. Meanwhile, to obtain the non-Brownian functionals, whose path integrals are over Lévy trajectories, the space-fractional Fokker-Planck equation and the tempered space fractional diffusion equations have been widely used; see [19, 22, 27].

The current work is devoted to proposing a second-order Grünwald-Letnikov-formula-based approximation for the fractional substantial derivative (1.1), and applying it to derive a high-order fully discrete scheme for the time fractional

substantial diffusion equation (TFSDE)

$${}_0D_t^{\alpha,\lambda}u(\mathbf{x},t) = \Delta u(\mathbf{x},t) + F(\mathbf{x},t), \quad \mathbf{x} \in \Omega, \quad t \in (0, T], \quad (1.2)$$

$$u(\mathbf{x}, 0) = u_0(\mathbf{x}), \quad \mathbf{x} \in \Omega, \quad (1.3)$$

$$u(\mathbf{x}, t)|_{\mathbf{x} \in \partial\Omega} = \phi(\mathbf{x}, t), \quad t \in (0, T], \quad (1.4)$$

where Δ is the Laplacian operator, \mathbf{x} denotes the one-dimensional or two-dimensional space variable, $\partial\Omega$ is the boundary of domain Ω , $F(\mathbf{x}, t)$, $u_0(\mathbf{x})$ and $\phi(\mathbf{x}, t)$ are given functions ; ${}_0D_t^{\alpha,\lambda}$ is the substantial derivative defined by (1.1), and $0 < \alpha \leq 1$.

The main novelty of our paper is the derivation of a second-order operator, which is based on the modified definition of the Grünwald derivative (see section 3.4, [29]), for the approximation of the fractional substantial derivative. The modified Grünwald derivative is defined by [29]

$${}_0^{GL}D_t^\alpha f(t) \equiv \lim_{\tau \rightarrow 0} \frac{1}{\tau^\alpha} \sum_{k=0}^{[t/\tau]} g_k^\alpha f(t - k\tau + \frac{\alpha}{2}\tau), \quad g_k^\alpha = (-1)^k \binom{\alpha}{k}. \quad (1.5)$$

Actually, if dropping the term $\frac{\alpha}{2}\tau$ off in the right side of above definition, one gets the original definition of the Grünwald derivative. The advantage of the modified Grünwald derivative is that it permits the design of more efficient algorithms to approximate the Riemann-Liouville fractional derivatives than using the shifted Grünwald-Letnikov formula directly. In this paper, by developing the modified Grünwald derivative and the shifted Grünwald-Letnikov formula to the fractional substantial derivative, a modified shifted substantial Grünwald formula and its asymptotic expansion are presented. Based upon the asymptotic expansion, a second-order approximation of the fractional substantial derivative is derived.

There are also some other approaches for the approximation of fractional derivatives, such as the $L1$ approximation [23, 34], the fractional linear multi-step methods (FLMMs) developed by Lubich [42], the $L2$ approximation with using superior convergence [1, 18], etc. However, **to the best of authors' knowledge**, very limited work has been presented for the fractional substantial derivative. Chen and Deng [11] extended the p -th order FLMMs [25] to approximate the fractional substantial derivative, and applied it to solve the fractional Feynman-Fac equation [14] lately. Very recently, Chen and Deng [12] proposed some algorithms for the equation with the fractional substantial derivative in time and the tempered fractional derivatives in space, in which the numerical stability and error estimate have been given for a scheme with **the** first-order accuracy in time and **the** second-order accuracy in space.

The main goal of our paper is to construct a second-order approximation for the time fractional substantial derivative, and subsequently to solve the TFSDE (1.2)-(1.4) by combining the existed fourth-order **compact approximation** for the space derivatives [36], and establish the numerical stability and error estimate of the derived fully discretized scheme.

In this work, we assume that the solution to the underlying equation satisfies suitable regularity requirements. The assumption can be satisfied in certain conditions, while it may not hold for many time-fractional differential equations; see related discussion for the case $\lambda = 0$ in [7, 20, 41]. To circumvent the requirement of high regularity of the solution, we apply starting quadrature to add correction terms in the proposed scheme. The starting quadrature was first developed in [25], and **has** been used to deal with problems with nonsmooth solution; see [14, 42]. The validity of the proposed algorithm is illustrated in Example 6.3 by solving a two-dimensional time fractional substantial diffusion equation.

The remainder of this paper is organized as follows. In Section 2, a second-order operator for the approximation of the fractional substantial derivative is derived. In Section 3, the proposed approximation is applied to TFSDE (1.2)-(1.4), and a fully discretized scheme is derived by combining the fourth-order compact formula in space. In Section 4, we give a discrete prior estimate for the numerical solution, and then prove the convergence and stability of the proposed scheme. The behavior of our proposed scheme when applied to solve problems with non-smooth solution is further discussed and the improved algorithm is proposed in Section 5. Numerical results are presented in Section 6. Some concluding remarks are included in the final section.

2. A second-order approximation for the fractional substantial derivative

In this section, we first present a second-order approximation for the fractional substantial derivative. For $0 < \alpha \leq 1$, we give the following identities from (1.1):

$${}_a I_t^{\alpha, \lambda} f(t) = e^{-\lambda t} {}_a I_t^{\alpha} [e^{\lambda t} f(t)], \quad {}_a D_t^{\alpha, \lambda} f(t) = e^{-\lambda t} {}_a D_t^{\alpha} [e^{\lambda t} f(t)]. \quad (2.1)$$

Moreover, if $f(a) = 0$, by the composition formula [30], it holds that

$${}_a I_t^{\alpha, \lambda} {}_a D_t^{\alpha, \lambda} f(t) = f(t). \quad (2.2)$$

Based on the definition of the modified Grünwald derivative [29] and the

shifted Grünwald- Letnikov formula [37], we define

$$A_{\tau,r}^{\alpha,\lambda} f(t) = \frac{1}{\tau^\alpha} \sum_{k=0}^{+\infty} e^{-\lambda(k-r)\tau} g_k^\alpha f(t - (k-r)\tau), \quad \forall \alpha > 0. \quad (2.3)$$

where τ is the step size, and $\{g_k^\alpha\}$ are the coefficients of the power series expansion of the function $(1-z)^\alpha$, i.e.,

$$(1-z)^\alpha = \sum_{k=0}^{+\infty} (-1)^k \binom{\alpha}{k} z^k = \sum_{k=0}^{+\infty} g_k^\alpha z^k, \quad \forall |z| < 1. \quad (2.4)$$

Obviously, $g_k^\alpha = (-1)^k \binom{\alpha}{k}$ can also be evaluated recursively by

$$g_0^\alpha = 1, \quad g_k^\alpha = \left(1 - \frac{\alpha+1}{k}\right) g_{k-1}^\alpha, \quad k = 1, 2, \dots. \quad (2.5)$$

Let $f \in L_1(\mathbb{R})$. Define the function space $\mathcal{C}^{n+\alpha}(\mathbb{R})$ by

$$\mathcal{C}^{n+\alpha}(\mathbb{R}) = \left\{ f \mid \int_{-\infty}^{+\infty} (1+|\omega|)^{n+\alpha} |\hat{f}(\omega)| d\omega < \infty \right\},$$

where $\hat{f}(\omega) = \int_{-\infty}^{+\infty} e^{i\omega t} f(t) dt$ is the Fourier transform of $f(t)$.

Motivated by the work in [37], we present an asymptotic expansion of $A_{\tau,r}^{\alpha,\lambda} f(t)$.

Lemma 2.1. *Suppose that $f \in L_1(\mathbb{R})$ and $f \in \mathcal{C}^{n+\alpha}(\mathbb{R})$. Then*

$$A_{\tau,r}^{\alpha,\lambda} f(t) = -\infty D_t^{\alpha,\lambda} f(t) + \sum_{k=1}^{n-1} c_{r,k}^\alpha -\infty D_t^{\alpha+k,\lambda} f(t) \tau^k + \mathcal{O}(\tau^n) \quad (2.6)$$

uniformly holds in $t \in \mathbb{R}$ as $\tau \rightarrow 0$, where $c_{r,k}^\alpha$ are the coefficients of the power series expansion of the function $W_{\alpha,r}(z) = \left(\frac{1-e^{-z}}{z}\right)^\alpha e^{rz}$.

Proof Using the Fourier transform, we obtain

$$\begin{aligned} \mathcal{F}(A_{\tau,r}^{\alpha,\lambda} f)(\omega) &= \frac{1}{\tau^\alpha} \sum_{k=0}^{\infty} e^{-\lambda(k-r)\tau} g_k^\alpha \mathcal{F}(f(t - (k-r)\tau))(\omega) \\ &= \frac{1}{\tau^\alpha} \sum_{k=0}^{\infty} e^{-\lambda(k-r)\tau} g_k^\alpha e^{i\omega(k-r)\tau} \hat{f}(\omega) \\ &= \frac{1}{\tau^\alpha} (1 - e^{-(\lambda-i\omega)\tau})^\alpha e^{r(\lambda-i\omega)\tau} \hat{f}(\omega) \\ &= (\lambda - i\omega)^\alpha \left(\frac{1 - e^{-(\lambda-i\omega)\tau}}{(\lambda - i\omega)\tau} \right)^\alpha e^{r(\lambda-i\omega)\tau} \hat{f}(\omega) \\ &= (\lambda - i\omega)^\alpha W_{\alpha,r}((\lambda - i\omega)\tau) \hat{f}(\omega). \end{aligned}$$

Observe that $c_{r,0}^\alpha = 1$ and

$$\mathcal{F}(-_\infty D_t^{\alpha+k,\lambda} f)(\omega) = (\lambda - i\omega)^{\alpha+k} \hat{f}(\omega)$$

for $k = 0, 1, 2 \dots$. Therefore, there exists

$$\mathcal{F}(A_{\tau,r}^{\alpha,\lambda} f)(\omega) = \sum_{k=0}^{n-1} c_{r,k}^\alpha \mathcal{F}(-_\infty D_t^{\alpha+k,\lambda} f)(\omega) \tau^k + \Phi(\omega, \tau), \quad (2.7)$$

where

$$\Phi(\omega, \tau) = (\lambda - i\omega)^\alpha [W_{\alpha,r}((\lambda - i\omega)\tau) - \sum_{k=0}^{n-1} c_{r,k}^\alpha (\lambda - i\omega)^k \tau^k] \hat{f}(\omega).$$

Taking inverse Fourier **transform** on the both sides of (2.7) leads to

$$A_{\tau,r}^{\alpha,\lambda} f(t) - \sum_{k=0}^{n-1} c_{r,k}^\alpha -_\infty D_t^{\alpha+k,\lambda} f(t) \tau^k = \frac{1}{2\pi} \int_{-\infty}^{+\infty} \Phi(\omega, \tau) e^{-i\omega t} d\omega. \quad (2.8)$$

Due to the fact that $f \in \mathcal{C}^{n+\alpha}(\mathbb{R})$ and $W_{\alpha,r}(z) = \sum_{k=0}^{n-1} c_{r,k}^\alpha z^k + O(z^n)$, then there exists a constant c such that

$$\begin{aligned} & A_{\tau,r}^{\alpha,\lambda} f(t) - -_\infty D_t^{\alpha,\lambda} f(t) - \sum_{k=1}^{n-1} c_{r,k}^\alpha -_\infty D_t^{\alpha+k,\lambda} f(t) \tau^k \\ & \leq \frac{1}{2\pi} \int_{-\infty}^{+\infty} |\Phi(\omega, \tau)| d\omega \leq c \int_{-\infty}^{+\infty} (1 + |\omega|)^{n+\alpha} |\hat{f}(\omega)| d\omega \cdot \tau^n = \mathcal{O}(\tau^n). \end{aligned}$$

for sufficiently small τ . This ends the proof. \square

Remark 2.2. Take $\lambda = 0$, $\alpha = 1$ in above lemma. When $r = 0$, (2.6) reduces to the backward difference operator for the standard derivative:

$$f'(t) = \frac{f(t) - f(t - \tau)}{\tau} + \mathcal{O}(\tau);$$

when $r = \frac{\alpha}{2} = \frac{1}{2}$, (2.6) reduces to the central difference quotient operator:

$$f'(t) = \frac{f(t + \frac{1}{2}\tau) - f(t - \frac{1}{2}\tau)}{\tau} + \mathcal{O}(\tau^2).$$

On the other hand, the accuracy can be improved by shifting from the backward difference formula to the central difference counterpart. From this point of view, the fractional shift in the formula (2.6) is natural and reasonable to achieve high accuracy.

From Lemma 2.1, we can take $A_{\tau,r}^{\alpha,\lambda} f(t)$ as a first-order approximation to the fractional substantial derivative ${}_{-\infty}D_t^{\alpha,\lambda} f(t)$, that is,

$$A_{\tau,r}^{\alpha,\lambda} f(t) = {}_{-\infty}D_t^{\alpha,\lambda} f(t) + \mathcal{O}(\tau),$$

which is consistent with the approximation of the Riemann-Liouville derivative by the shifted Grünwald-Letnikov formula [37]. Note that $c_{r,1}^\alpha = r - \frac{\alpha}{2}$. We take $t = s - r\tau$ and $r = \frac{\alpha}{2}$ in (2.6), then it follows that

$$A_{\tau,r}^{\alpha,\lambda} f(s - \frac{\alpha}{2}\tau) = {}_{-\infty}D_t^{\alpha,\lambda} f(s - \frac{\alpha}{2}\tau) + \mathcal{O}(\tau^2).$$

By (2.3), and using the Lagrange linear interpolation at the points $t = s$ and $t = s - \tau$ for the approximation of ${}_{-\infty}D_t^{\alpha,\lambda} f(s - r\tau)$, and replacing s with t , we can derive the following second-order Grünwald-Letnikov-formula-based approximation for the fractional substantial derivative.

Theorem 2.3. *Suppose that $f \in L_1(\mathbb{R})$ and $f \in \mathcal{C}^{2+\alpha}(\mathbb{R})$. Then*

$$(1 - \frac{\alpha}{2}) {}_{-\infty}D_t^{\alpha,\lambda} f(t) + \frac{\alpha}{2} {}_{-\infty}D_t^{\alpha,\lambda} f(t - \tau) = \frac{1}{\tau^\alpha} \sum_{k=0}^{+\infty} g_k^{\alpha,\lambda} f(t - k\tau) + \mathcal{O}(\tau^2)$$

uniformly holds in $t \in \mathbb{R}$ as $\tau \rightarrow 0$, where $g_k^{\alpha,\lambda} = e^{-(k-\frac{\alpha}{2})\lambda\tau} g_k^\alpha$.

To apply Theorem 2.3 to the generic function $f(t)$ ($f(0) = 0$) on $[0, +\infty)$, we make the zero extension as follows:

$$\tilde{f}(t) = \begin{cases} f(t), & t \in [0, +\infty), \\ 0, & \text{others} \end{cases}$$

and further assume that $\tilde{f}(t)$ satisfies the assumptions in Theorem 2.3, i.e., $\tilde{f}(t) \in \mathcal{C}^{2+\alpha}(\mathbb{R})$. Then it follows that

$$(1 - \frac{\alpha}{2}) {}_0D_t^{\alpha,\lambda} f(t) + \frac{\alpha}{2} {}_0D_t^{\alpha,\lambda} f(t - \tau) = \frac{1}{\tau^\alpha} \sum_{k=0}^{[\frac{t}{\tau}]} g_k^{\alpha,\lambda} f(t - k\tau) + \mathcal{O}(\tau^2). \quad (2.9)$$

3. Derivation of the numerical scheme

In this section, we construct a numerical scheme for the problem (1.2)-(1.4). We first derive a semi-discretized scheme by the proposed approximation (2.9). Subsequently, with the use of the fourth-order compact finite difference approximation of the space derivatives [36], we give a fully discrete scheme. In what follows, we suppose that $u(\mathbf{x}, 0) = 0$ without loss of generality. Otherwise, consider the equation with the solution $v(\mathbf{x}, t) = u(\mathbf{x}, t) - e^{-\lambda t} u(\mathbf{x}, 0)$ instead. Furthermore, assume that the extended $u(\cdot, t) \in \mathcal{C}^{2+\alpha}(\mathbb{R})$.

3.1. A semi-discrete scheme

Now we are in the position to discrete (1.2) in time. Take the uniform time step size $\tau = T/N$ and let $t_n = n\tau$, $0 \leq n \leq N$. The temporal domain $[0, T]$ is covered by $\Omega_\tau = \{t_n \mid 0 \leq n \leq N\}$. For a given grid function $v = \{v^n\}_{n=0}^N$ defined on Ω_τ , we introduce the average operator:

$$\mathcal{A}_t^\alpha v^n = (1 - \frac{\alpha}{2})v^n + \frac{\alpha}{2}v^{n-1} \quad (3.1)$$

and the discrete difference quotient operator:

$$\delta_t^{\alpha, \lambda} v^n = \frac{1}{\tau^\alpha} \sum_{k=0}^n g_k^{\alpha, \lambda} v^{n-k}. \quad (3.2)$$

Considering the equation (1.2) at the grid points (\mathbf{x}, t_n) , we have

$${}_0D_t^{\alpha, \lambda} u(\mathbf{x}, t_n) = \Delta u(\mathbf{x}, t_n) + F(\mathbf{x}, t_n), \quad 0 \leq n \leq N. \quad (3.3)$$

Performing the average operator \mathcal{A}_t^α defined in (3.1) on the equation (3.3), we have

$$\mathcal{A}_t^\alpha {}_0D_t^{\alpha, \lambda} u(\mathbf{x}, t_n) = \mathcal{A}_t^\alpha \Delta u(\mathbf{x}, t_n) + \mathcal{A}_t^\alpha F(\mathbf{x}, t_n), \quad 1 \leq n \leq N.$$

Applying the formula (2.9) and combining (3.2) yield

$$\delta_t^{\alpha, \lambda} u(\mathbf{x}, t_n) = \mathcal{A}_t^\alpha \Delta u(\mathbf{x}, t_n) + \mathcal{A}_t^\alpha F(\mathbf{x}, t_n) + \mathcal{O}(\tau^2), \quad 1 \leq n \leq N. \quad (3.4)$$

Noting that (3.4) is a semi-discretization to (1.2), in next subsection, we will derive a fully discretization by the finite difference method. It is worth to point out that (3.4) can also be solved by many other methods that we mentioned in the introduction part, such as spectral method, finite element method, **discontinuous Galerkin method**, etc.

3.2. A fully discrete scheme

In this part, we use the fourth-order compact **approximation** for the spatial discretization of the problem (1.2)-(1.4). To fix the idea, we consider only one-dimensional problem (1.2)-(1.4) in this subsection and illustrate the **generalization to** two-dimensional case via an example in Section 6.

Let $\Omega = (a, b)$. Take a positive integer M and let $h = (b - a)/M$, $x_i = ih$, $0 \leq i \leq M$. Then the spatial domain $[a, b]$ is covered by $\Omega_h = \{x_i \mid 0 \leq i \leq M\}$.

For completeness of the paper, we introduce some results that are helpful for understanding the construction of the scheme. The following lemma is required to achieve high-order accuracy for spatial discretization.

Lemma 3.1. [36] Denote $\theta(s) = (1-s)^3[5-3(1-s)^2]$. If $G(x) \in C^6[x_{i-1}, x_{i+1}]$, $x_{i+1} = x_i + h$, $x_{i-1} = x_i - h$, it holds that

$$\begin{aligned} \frac{1}{12}[G''(x_{i-1}) + 10G''(x_i) + G''(x_{i+1})] &= \frac{G(x_{i-1}) - 2G(x_i) + G(x_{i+1}))}{h^2} \\ &+ \frac{h^4}{360} \int_0^1 [G^{(6)}(x_i - sh) + G^{(6)}(x_i + sh)]\theta(s)ds, \quad 1 \leq i \leq M-1. \end{aligned}$$

Introduce the average operator \mathcal{A}_x in spatial direction as

$$\mathcal{A}_x v_i = \begin{cases} \frac{1}{12}(v_{i-1} + 10v_i + v_{i+1}), & 1 \leq i \leq M-1, \\ v_i, & i = 0 \text{ or } M. \end{cases}$$

Denote by δ_x^2 the second-order central difference quotient operator, that is

$$\delta_x^2 v_i = \frac{1}{h^2}(v_{i-1} - 2v_i + v_{i+1}).$$

Taking $\mathbf{x} = x_i$ and performing the average operator \mathcal{A}_x on both sides of (3.4) gives

$$\begin{aligned} \mathcal{A}_x \delta_t^{\alpha, \lambda} u(x_i, t_n) &= \mathcal{A}_x \mathcal{A}_t^\alpha \partial_x^2 u(x_i, t_n) + \mathcal{A}_x \mathcal{A}_t^\alpha \mathbf{F}(x_i, t_n) + \mathcal{O}(\tau^2), \\ 1 \leq i \leq M-1, \quad 1 \leq n \leq N. \end{aligned} \quad (3.5)$$

Define the grid functions

$$U_i^n = u(x_i, t_n), \quad F_i^n = F(x_i, t_n), \quad 0 \leq i \leq M, \quad 0 \leq n \leq N.$$

Note that the operators \mathcal{A}_t^α and δ_x^2 can be commuted to each other. Then using Lemma 3.1 yields

$$\mathcal{A}_x \delta_t^{\alpha, \lambda} U_i^n - \mathcal{A}_t^\alpha \delta_x^2 U_i^n = \mathcal{A}_t^\alpha \mathcal{A}_x F_i^n + R_i^n, \quad 1 \leq i \leq M-1, \quad 1 \leq n \leq N, \quad (3.6)$$

where there exists a constant c_1 such that

$$|R_i^n| \leq c_1(\tau^2 + h^4), \quad 1 \leq i \leq M-1, \quad 1 \leq n \leq N. \quad (3.7)$$

Omitting the small terms R_i^n in (3.6), denoting u_i^n as the approximation of U_i^n and noticing the initial and boundary conditions

$$U_i^0 = 0, \quad 0 \leq i \leq M, \quad (3.8)$$

$$U_0^n = \phi(x_0, \mathbf{t}_n), \quad U_M^n = \phi(x_M, \mathbf{t}_n), \quad 1 \leq n \leq N, \quad (3.9)$$

we obtain a (α, λ) -dependent compact difference scheme

$$\mathcal{A}_x \delta_t^{\alpha, \lambda} u_i^n - \mathcal{A}_t^\alpha \delta_x^2 u_i^n = \mathcal{A}_t^\alpha \mathcal{A}_x F_i^n, \quad 1 \leq i \leq M-1, \quad 1 \leq n \leq N, \quad (3.10)$$

$$u_i^0 = 0, \quad 0 \leq i \leq M, \quad (3.11)$$

$$u_0^n = \phi(x_0, \mathbf{t}_n), \quad u_M^n = \phi(x_M, \mathbf{t}_n), \quad 1 \leq n \leq N. \quad (3.12)$$

Remark 3.2. When $\lambda = \lambda(x)$, the problem (1.2)-(1.4) becomes the Feynman-Kac equation considered in [14]. In Section 5, we will show numerically that the adapted scheme, i.e., taking $\lambda_i = \lambda(x_i)$ in above scheme, is of second-order convergence in time and fourth-order convergence in space for the forward Feynman-Kac equation considered in [14].

4. Stability and convergence analysis

In this section, we shall give the stability and convergence analysis for the scheme (3.10)-(3.12).

Let $V_h = \{v \mid v = (v_0, \dots, v_M) \text{ is a grid function on } \Omega_h \text{ and } v_0 = v_M = 0\}$. For any $u, v \in V_h$, we denote the maximum norm by $\|u\|_\infty = \max_{0 \leq i \leq M} |u_i|$ and define the discrete inner products and induced norm as

$$(u, v) = h \sum_{i=1}^{M-1} u_i v_i, \quad \|u\| = \sqrt{(u, u)}.$$

Let

$$l_n^\alpha = \sum_{k=0}^n g_k^\alpha, \quad n \geq 0. \quad (4.1)$$

Then l_n^α are the coefficients of the power series expansion of the function $(1 - z)^{\alpha-1}$, i.e.,

$$(1 - z)^{\alpha-1} = \sum_{k=0}^{+\infty} l_k^\alpha z^k,$$

In fact, from $(1 - z)^\alpha = (1 - z) \cdot (1 - z)^{\alpha-1}$, we have

$$\sum_{k=0}^{+\infty} g_k^\alpha z^k = (1 - z) \sum_{k=0}^{+\infty} l_k^\alpha z^k = l_0^\alpha + \sum_{k=1}^{+\infty} (l_k^\alpha - l_{k-1}^\alpha) z^k.$$

Comparing the coefficients of power series expansion on both sides leads to the desired result. In addition, the coefficients l_n^α in (4.1) satisfy the following

properties for $0 < \alpha < 1$,

$$\begin{cases} l_0^\alpha = 1, & l_1^\alpha = 1 - \alpha, \\ l_0^\alpha \geq l_1^\alpha \geq l_2^\alpha \geq l_3^\alpha \geq \dots \geq 0. \end{cases} \quad (4.2)$$

And it has the lower bound, i.e.,

$$l_{n-1}^\alpha \geq \frac{1}{n^\alpha \Gamma(1 - \alpha)}. \quad (4.3)$$

See Lemma 2.1 in [14] for details.

Prior to **considering** the stability and convergence of the scheme (3.10)-(3.12), we need to provide some auxiliary lemmas. Inspired by the work in [1], we first give the following lemma.

Lemma 4.1. *For a given sequence $\{v^n\}_{n=0}^\infty$, we have*

$$v^n \delta_t^{\alpha, \lambda} v^n \geq \frac{1}{2} e^{-\frac{\alpha}{2} \lambda \tau} \delta_t^{\alpha, 2\lambda} (v^n)^2 + \frac{e^{-\frac{\alpha}{2} \lambda \tau}}{2 l_0^\alpha} \tau^\alpha (\delta_t^{\alpha, \lambda} v^n)^2, \quad (4.4)$$

$$v^{n-1} \delta_t^{\alpha, \lambda} v^n \geq \frac{1}{2} e^{(1-\frac{\alpha}{2}) \lambda \tau} \delta_t^{\alpha, 2\lambda} (v^n)^2 - \frac{e^{(1-\frac{\alpha}{2}) \lambda \tau}}{2(l_0^\alpha - l_1^\alpha)} \tau^\alpha (\delta_t^{\alpha, \lambda} v^n)^2, \quad (4.5)$$

where

$$\delta_t^{\alpha, 2\lambda} (v^n)^2 = \frac{1}{\tau^\alpha} \sum_{k=0}^n g_k^{\alpha, 2\lambda} (v^{n-k})^2.$$

Proof Note that $g_k^{\alpha, \lambda} = e^{\lambda(\frac{\alpha}{2}-k)\tau} g_k^\alpha$. The expressions $\delta_t^{\alpha, \lambda} v^n$, $\delta_t^{\alpha, 2\lambda} (v^n)^2$ can also be rewritten as

$$\delta_t^{\alpha, \lambda} v^n = e^{\lambda(\frac{\alpha}{2}-n)\tau} \frac{1}{\tau^\alpha} \sum_{k=0}^n l_{n-k}^\alpha (e^{\lambda k \tau} v^k - e^{\lambda(k-1)\tau} v^{k-1}), \quad (4.6)$$

$$\delta_t^{\alpha, 2\lambda} (v^n)^2 = e^{2\lambda(\frac{\alpha}{2}-n)\tau} \frac{1}{\tau^\alpha} \sum_{k=0}^n l_{n-k}^\alpha [(e^{\lambda k \tau} v^k)^2 - (e^{\lambda(k-1)\tau} v^{k-1})^2], \quad (4.7)$$

where $v^{-1} = 0$. To **simplify** notation but without ambiguity, hereafter, we denote

$v_\lambda^k := e^{\lambda k \tau} v^k$. Thus we have

$$\begin{aligned}
& e^{(2n-\frac{\alpha}{2})\lambda\tau} [v^n \delta_t^{\alpha,\lambda} v^n - \frac{1}{2} e^{-\frac{\alpha}{2}\lambda\tau} \delta_t^{\alpha,2\lambda} (v^n)^2] \\
&= v_\lambda^n \frac{1}{\tau^\alpha} \sum_{k=0}^n l_{n-k}^\alpha (v_\lambda^k - v_\lambda^{k-1}) - \frac{1}{\tau^\alpha} \sum_{k=0}^n l_{n-k}^\alpha (v_\lambda^k - v_\lambda^{k-1}) \left(\frac{v_\lambda^k + v_\lambda^{k-1}}{2} \right) \\
&= \frac{1}{\tau^\alpha} \sum_{k=0}^n l_{n-k}^\alpha (v_\lambda^k - v_\lambda^{k-1}) \left(v_\lambda^n - \frac{v_\lambda^k + v_\lambda^{k-1}}{2} \right) \\
&= \frac{1}{\tau^\alpha} \sum_{k=0}^n l_{n-k}^\alpha (v_\lambda^k - v_\lambda^{k-1}) \left[\frac{v_\lambda^k - v_\lambda^{k-1}}{2} + \sum_{j=k+1}^n (v_\lambda^j - v_\lambda^{j-1}) \right] \\
&= \frac{1}{2\tau^\alpha} \sum_{k=0}^n l_{n-k}^\alpha (v_\lambda^k - v_\lambda^{k-1})^2 + \frac{1}{\tau^\alpha} \sum_{j=1}^n (v_\lambda^j - v_\lambda^{j-1}) \sum_{k=0}^{j-1} l_{n-k}^\alpha (v_\lambda^k - v_\lambda^{k-1}).
\end{aligned} \tag{4.8}$$

Denote

$$w_\lambda^j = \sum_{k=0}^{j-1} l_{n-k}^\alpha (v_\lambda^k - v_\lambda^{k-1}), \quad 1 \leq j \leq n \tag{4.9}$$

and

$$w_\lambda^{n+1} = \sum_{k=0}^n l_{n-k}^\alpha (v_\lambda^k - v_\lambda^{k-1}) = \tau^\alpha e^{(n-\frac{\alpha}{2})\lambda\tau} \delta_t^{\alpha,\lambda} v^n. \tag{4.10}$$

Then it follows that

$$v_\lambda^0 - v_\lambda^{-1} = \frac{w_\lambda^1}{l_n^\alpha}; \quad v_\lambda^j - v_\lambda^{j-1} = \frac{w_\lambda^{j+1} - w_\lambda^j}{l_{n-j}^\alpha}, \quad 1 \leq j \leq n. \tag{4.11}$$

Substituting (4.9)-(4.11) into (4.8) leads to

$$\begin{aligned}
& e^{(2n-\frac{\alpha}{2})\lambda\tau} [v^n \delta_t^{\alpha,\lambda} v^n - \frac{1}{2} e^{-\frac{\alpha}{2}\lambda\tau} \delta_t^{\alpha,2\lambda} (v^n)^2] \\
&= \frac{1}{2\tau^\alpha} l_n^\alpha \left(\frac{w_\lambda^1}{l_n^\alpha} \right)^2 + \frac{1}{2\tau^\alpha} \sum_{k=1}^n l_{n-k}^\alpha \left(\frac{w_\lambda^{k+1} - w_\lambda^k}{l_{n-k}^\alpha} \right)^2 + \frac{1}{\tau^\alpha} \sum_{j=1}^n \left(\frac{w_\lambda^{j+1} - w_\lambda^j}{l_{n-j}^\alpha} \right) w_\lambda^j \\
&= \frac{1}{2\tau^\alpha} \frac{(w_\lambda^{n+1})^2}{l_0^\alpha} + \frac{1}{2\tau^\alpha} \sum_{k=1}^n \left(\frac{1}{l_{n+1-k}^\alpha} - \frac{1}{l_{n-k}^\alpha} \right) (w_\lambda^k)^2.
\end{aligned} \tag{4.12}$$

Multiplying (4.12) by $e^{(\frac{\alpha}{2}-2n)\lambda\tau}$ and recalling (4.2), we have

$$v^n \delta_t^{\alpha,\lambda} v^n - \frac{1}{2} e^{-\frac{\alpha}{2}\lambda\tau} \delta_t^{\alpha,2\lambda} (v^n)^2 \geq \frac{e^{(\frac{\alpha}{2}-2n)\lambda\tau}}{2\tau^\alpha} \frac{(w_\lambda^{n+1})^2}{l_0^\alpha} = \frac{e^{-\frac{\alpha}{2}\lambda\tau}}{2l_0^\alpha} \tau^\alpha (\delta_t^{\alpha,\lambda} v^n)^2.$$

Next, we prove the inequality (4.5). Considering the difference, using the notations (4.9)-(4.10) again and noting the equality (4.12), we have

$$\begin{aligned}
& e^{(2n-1-\frac{\alpha}{2})\lambda\tau} [v^{n-1}\delta_t^{\alpha,\lambda}v^n - \frac{1}{2}e^{(1-\frac{\alpha}{2})\lambda\tau}\delta_t^{\alpha,2\lambda}(v^n)^2 + \frac{e^{(1-\frac{\alpha}{2})\lambda\tau}}{2(l_0^\alpha - l_1^\alpha)}\tau^\alpha(\delta_t^{\alpha,\lambda}v^n)^2] \\
&= e^{(2n-\frac{\alpha}{2})\lambda\tau} [v^n\delta_t^{\alpha,\lambda}v^n - \frac{1}{2}e^{-\frac{\alpha}{2}\lambda\tau}\delta_t^{\alpha,2\lambda}(v^n)^2] + \frac{\tau^\alpha}{2(l_0^\alpha - l_1^\alpha)}e^{(2n-\alpha)\lambda\tau}(\delta_t^{\alpha,\lambda}v^n)^2 \\
&\quad - [e^{n\lambda\tau}v^n - e^{(n-1)\lambda\tau}v^{n-1}]e^{(n-\frac{\alpha}{2})\lambda\tau}\delta_t^{\alpha,\lambda}v^n \\
&= \frac{1}{2\tau^\alpha}\frac{(w_\lambda^{n+1})^2}{l_0^\alpha} + \frac{1}{2\tau^\alpha}\sum_{k=1}^n\left(\frac{1}{l_{n+1-k}^\alpha} - \frac{1}{l_{n-k}^\alpha}\right)(w_\lambda^k)^2 + \frac{\tau^\alpha}{2(l_0^\alpha - l_1^\alpha)}\left(\frac{w_\lambda^{n+1}}{\tau^\alpha}\right)^2 \\
&\quad - \frac{(w_\lambda^{n+1} - w_\lambda^n)w_\lambda^{n+1}}{l_0^\alpha\tau^\alpha} \\
&= \frac{1}{\tau^\alpha}\left(\frac{1}{2(l_0^\alpha - l_1^\alpha)} - \frac{1}{2l_0^\alpha}\right)(w_\lambda^{n+1})^2 + \frac{1}{\tau^\alpha}\frac{1}{l_0^\alpha}w_\lambda^n w_\lambda^{n+1} + \frac{1}{2\tau^\alpha}\left(\frac{1}{l_1^\alpha} - \frac{1}{l_0^\alpha}\right)(w_\lambda^n)^2 \\
&\quad + \frac{1}{2\tau^\alpha}\sum_{k=1}^{n-1}\left(\frac{1}{l_{n+1-k}^\alpha} - \frac{1}{l_{n-k}^\alpha}\right)(w_\lambda^k)^2 \\
&= \frac{1}{2l_0^\alpha\tau^\alpha}\left(\sqrt{\frac{l_1^\alpha}{l_0^\alpha - l_1^\alpha}}w_\lambda^{n+1} + \sqrt{\frac{l_0^\alpha - l_1^\alpha}{l_1^\alpha}}w_\lambda^n\right)^2 + \frac{1}{2\tau^\alpha}\sum_{k=1}^{n-1}\left(\frac{1}{l_{n+1-k}^\alpha} - \frac{1}{l_{n-k}^\alpha}\right)(w_\lambda^k)^2 \\
&\geq 0,
\end{aligned}$$

which completes the proof. \square

Lemma 4.2. *Given a sequence $\{v^n\}$ we have*

$$\begin{aligned}
\mathcal{A}_t^\alpha v^n \cdot \delta_t^{\alpha,\lambda}v^n &\geq \frac{1}{2}\left[\frac{\alpha}{2}e^{\lambda\tau} + \left(1 - \frac{\alpha}{2}\right)\right]e^{-\frac{\alpha}{2}\lambda\tau}\delta_t^{\alpha,2\lambda}(v^n)^2 \\
&\quad + \frac{2 - \alpha - e^{\lambda\tau}}{4}\tau^\alpha e^{-\frac{\alpha}{2}\lambda\tau}(\delta_t^{\alpha,\lambda}v^n)^2.
\end{aligned} \tag{4.13}$$

In particular, when $\alpha = 1$, $\lambda = 0$ it holds that

$$\mathcal{A}_t v^n \cdot \delta_t v^n := \frac{v^n + v^{n-1}}{2} \cdot \frac{v^n - v^{n-1}}{\tau} = \frac{(v^n)^2 - (v^{n-1})^2}{2\tau}.$$

Proof From Lemma 4.1, we have

$$\begin{aligned}
\mathcal{A}_t^\alpha v^n \cdot \delta_t^{\alpha, \lambda} v^n &= (1 - \frac{\alpha}{2}) v^n \delta_t^{\alpha, \lambda} v^n + \frac{\alpha}{2} v^{n-1} \delta_t^{\alpha, \lambda} v^n \\
&\geq (1 - \frac{\alpha}{2}) e^{-\frac{\alpha}{2} \lambda \tau} \left[\frac{1}{2} \delta_t^{\alpha, 2\lambda} (v^n)^2 + \frac{\tau^\alpha}{2l_0^\alpha} (\delta_t^{\alpha, \lambda} v^n)^2 \right] \\
&\quad + \frac{\alpha}{2} e^{(1-\frac{\alpha}{2})\lambda\tau} \left[\frac{1}{2} \delta_t^{\alpha, 2\lambda} (v^n)^2 - \frac{\tau^\alpha}{2(l_0^\alpha - l_1^\alpha)} (\delta_t^{\alpha, \lambda} v^n)^2 \right] \\
&= \frac{1}{2} \left[\frac{\alpha}{2} e^{\lambda\tau} + (1 - \frac{\alpha}{2}) \right] e^{-\frac{\alpha}{2} \lambda \tau} \delta_t^{\alpha, 2\lambda} (v^n)^2 \\
&\quad + \left[(1 - \frac{\alpha}{2}) \frac{1}{2l_0^\alpha} - \frac{\alpha}{2} \frac{e^{\lambda\tau}}{2(l_0^\alpha - l_1^\alpha)} \right] \tau^\alpha e^{-\frac{\alpha}{2} \lambda \tau} (\delta_t^{\alpha, \lambda} v^n)^2.
\end{aligned}$$

Substituting $l_0^\alpha = 1$ and $l_1^\alpha = 1 - \alpha$ into above inequality leads to (4.13), which completes the proof. \square

Lemma 4.3. [35] For any $u, v \in V_h$, it holds that $(\mathcal{A}_x u, v) = (u, \mathcal{A}_x v)$.

Lemma 4.4. [35] For any $v \in V_h$, it holds that

$$\frac{2}{3} \|v\|^2 \leq (\mathcal{A}_x v, v) \leq \|v\|^2.$$

Since \mathcal{A}_x is positive definite and self-adjoint, we can consider its square root denoted as Q_x , and have

$$(\mathcal{A}_x u, v) = (Q_x u, Q_x v). \quad (4.14)$$

Naturally, we define an equivalent norm as

$$\|v\|_A = \sqrt{(\mathcal{A}_x v, v)} = \sqrt{(Q_x v, Q_x v)}.$$

Lemma 4.5. [35] For any $v \in V_h$, it holds that $-(\delta_x^2 v, v) \geq C_\Omega \|v\|^2$, where C_Ω is a constant independent to the step size h but related to the domain Ω .

Now we give a prior estimate for the difference scheme (3.10)-(3.12).

Theorem 4.6. (Priori estimate) Suppose $\{v_i^n\}$ be the solution of

$$\mathcal{A}_x \delta_t^{\alpha, \lambda} v_i^n - \mathcal{A}_t^\alpha \delta_x^2 v_i^n = S_i^n, \quad 1 \leq i \leq M-1, \quad 1 \leq n \leq N, \quad (4.15)$$

$$v_0^n = 0, \quad v_M^n = 0, \quad 1 \leq n \leq N, \quad (4.16)$$

$$v_i^0 = v_0(x_i), \quad 0 \leq i \leq M. \quad (4.17)$$

If $2 - \alpha - e^{\lambda\tau} \geq 0$, then

$$\|v^n\|^2 \leq \frac{3}{2}e^{2|\lambda|T}\|v^0\|^2 + \frac{3\Gamma(1-\alpha)T^\alpha e^{2|\lambda|T}}{C_\Omega} \max_{0 \leq n \leq N} \|S^n\|^2 \quad (4.18)$$

for $1 < n \leq N$, where $v_0(x_0) = v_0(x_M) = 0$, $\|S^n\| = \sqrt{h \sum_{i=1}^{M-1} (S_i^n)^2}$.

Proof Taking the inner product of (4.15) with $\mathcal{A}_t^\alpha v^n$, we have

$$(\mathcal{A}_x \delta_t^{\alpha,\lambda} v^n, \mathcal{A}_t^\alpha v^n) - (\mathcal{A}_t^\alpha \delta_x^2 v^n, \mathcal{A}_t^\alpha v^n) = (S^n, \mathcal{A}_t^\alpha v^n). \quad (4.19)$$

By (4.14), for the first term on the left-hand side of (4.19), we get

$$(\mathcal{A}_x \delta_t^{\alpha,\lambda} v^n, \mathcal{A}_t^\alpha v^n) = (Q_x \delta_t^{\alpha,\lambda} v^n, Q_x \mathcal{A}_t^\alpha v^n) = (\delta_t^{\alpha,\lambda} Q_x v^n, \mathcal{A}_t^\alpha Q_x v^n).$$

By Lemma 4.2 and **assumption 2** $2 - \alpha - e^{\lambda\tau} \geq 0$, we have

$$\begin{aligned} (\delta_t^{\alpha,\lambda} Q_x v^n, \mathcal{A}_t^\alpha Q_x v^n) &= h \sum_{i=1}^{M-1} \delta_t^{\alpha,\lambda} (Q_x v^n)_i \cdot \mathcal{A}_t^\alpha (Q_x v^n)_i \\ &\geq \frac{1}{2} e^{-\frac{\alpha}{2}\lambda\tau} \left[\frac{\alpha}{2} e^{\lambda\tau} + \left(1 - \frac{\alpha}{2}\right) \right] h \sum_{i=1}^{M-1} \delta_t^{\alpha,2\lambda} (Q_x v^n)_i^2. \end{aligned} \quad (4.20)$$

For the second term on the left-hand side, from Lemma 4.5, we obtain

$$-(\mathcal{A}_t^\alpha \delta_x^2 v^n, \mathcal{A}_t^\alpha v^n) = -(\delta_x^2 \mathcal{A}_t^\alpha v^n, \mathcal{A}_t^\alpha v^n) \geq C_\Omega \|\mathcal{A}_t^\alpha v^n\|^2. \quad (4.21)$$

As to the term on the right-hand side, in view of Lemma 4.3 we have

$$(S^n, \mathcal{A}_t^\alpha v^n) \leq \|S^n\| \cdot \|\mathcal{A}_t^\alpha v^n\| \leq \frac{1}{4C_\Omega} \|S^n\|^2 + C_\Omega \|\mathcal{A}_t^\alpha v^n\|^2. \quad (4.22)$$

Substituting (4.20)-(4.22) into (4.19), we obtain

$$\frac{1}{2} e^{-\frac{\alpha}{2}\lambda\tau} \left[\frac{\alpha}{2} e^{\lambda\tau} + \left(1 - \frac{\alpha}{2}\right) \right] h \sum_{i=1}^{M-1} \delta_t^{\alpha,2\lambda} (Q_x v^n)_i^2 \leq \frac{1}{4C_\Omega} \|S^n\|^2.$$

Since $\frac{\alpha}{2} e^{\lambda\tau} + \left(1 - \frac{\alpha}{2}\right) \geq \frac{1}{2}$, we have

$$e^{-\frac{\alpha}{2}\lambda\tau} h \sum_{i=1}^{M-1} \delta_t^{\alpha,2\lambda} (Q_x v^n)_i^2 \leq \frac{1}{C_\Omega} \|S^n\|^2.$$

Consequently, by (4.7), it follows that

$$\begin{aligned}
& l_0^\alpha \|v_\lambda^n\|_A^2 \\
& \leq \sum_{k=1}^{n-1} (l_{n-k-1}^\alpha - l_{n-k}^\alpha) \|v_\lambda^k\|_A^2 + (l_{n-1}^\alpha - l_n^\alpha) \|v_\lambda^0\|_A^2 + \frac{\tau^\alpha}{C_\Omega} e^{(2n-\frac{\alpha}{2})\lambda\tau} \|S^n\|^2 \\
& \leq \sum_{k=1}^{n-1} (l_{n-k-1}^\alpha - l_{n-k}^\alpha) \|v_\lambda^k\|_A^2 + l_{n-1}^\alpha (\|v_\lambda^0\|_A^2 + \frac{\tau^\alpha}{C_\Omega l_{n-1}^\alpha} e^{(2n-\frac{\alpha}{2})\lambda\tau} \|S^n\|^2) \\
& \leq \sum_{k=1}^{n-1} (l_{n-k-1}^\alpha - l_{n-k}^\alpha) \|v_\lambda^k\|_A^2 + l_{n-1}^\alpha (\|v_\lambda^0\|^2 + \frac{\Gamma(1-\alpha)T^\alpha}{C_\Omega} e^{(2n-\frac{\alpha}{2})\lambda\tau} \|S^n\|^2), \\
& \quad 1 \leq n \leq N,
\end{aligned} \tag{4.23}$$

where we have used the estimate (4.3) in the last inequality. Denote

$$E^N = \|v_\lambda^0\|^2 + \frac{\Gamma(1-\alpha)T^\alpha}{C_\Omega} \max_{1 \leq n \leq N} e^{(2n-\frac{\alpha}{2})\lambda\tau} \max_{1 \leq n \leq N} \|S^n\|^2.$$

The inequality (4.23) is simplified to

$$l_0^\alpha \|v_\lambda^n\|_A^2 \leq \sum_{k=1}^{n-1} (l_{n-k-1}^\alpha - l_{n-k}^\alpha) \|v_\lambda^k\|_A^2 + l_{n-1}^\alpha E^N, \quad 1 \leq n \leq N. \tag{4.24}$$

Next we prove the following inequality

$$\|v_\lambda^n\|_A^2 \leq E^N, \quad 1 \leq n \leq N. \tag{4.25}$$

by the mathematical induction method. Obviously, it follows from (4.24) that the equality (4.25) holds for $n = 1$. Suppose that (4.25) still holds for $n = 1, 2, \dots, m$. From (4.24) at $n = m + 1$, we have

$$\begin{aligned}
l_0^\alpha \|v_\lambda^{m+1}\|_A^2 & \leq \sum_{k=1}^m (l_{m-k}^\alpha - l_{m+1-k}^\alpha) \|v_\lambda^k\|_A^2 + l_m^\alpha E^N \\
& \leq \sum_{k=1}^m (l_{m-k}^\alpha - l_{m+1-k}^\alpha) E^N + l_m^\alpha E^N = l_0^\alpha E^N,
\end{aligned}$$

which leads to (4.25).

It follows from Lemma 4.4 that

$$\frac{2}{3} e^{2\lambda n\tau} \|v^n\|^2 = \frac{2}{3} \|v_\lambda^n\|_A^2 \leq \|v_\lambda^n\|_A^2 \leq E^N.$$

Thus

$$\begin{aligned}
\|v^n\|^2 &\leq \frac{3}{2}e^{-2\lambda n\tau} E^N \\
&\leq \frac{3}{2}e^{-2\lambda n\tau} \left(\|v^0\|^2 + \frac{\Gamma(1-\alpha)T^\alpha}{C_\Omega} \max_{1 \leq n \leq N} e^{(2n-\frac{\alpha}{2})\lambda\tau} \max_{1 \leq n \leq N} \|S^n\|^2 \right) \\
&\leq \frac{3}{2}e^{2|\lambda|T} \|v^0\|^2 + \frac{3\Gamma(1-\alpha)T^\alpha}{C_\Omega} e^{2|\lambda|T} \max_{1 \leq n \leq N} \|S^n\|^2, \quad 1 \leq n \leq N.
\end{aligned}$$

This completes the proof. \square

Using Theorem 4.6, it is readily to obtain the following result.

Theorem 4.7. (Stability) *The difference scheme (3.10)-(3.12) is unconditionally stable **with respect to** the initial value and right-hand term.*

We now turn to the convergence of the difference scheme (3.10)-(3.12).

Theorem 4.8. (Convergence) *Assume that $u(x, t) \in C^{6,3}(\Omega \times [0, T])$ be the solution of the problem (1.2)-(1.4), and $\{u_i^n\}$ be the solution of difference scheme (3.10)-(3.12). We further assume that zero extended function $u(x, t)$ satisfies the assumptions in Theorem 2.3. Let $e_i^n = u(x_i, t_n) - u_i^n$, $0 \leq i \leq M$, $0 \leq n \leq N$. Then the following estimate*

$$\|e^n\| \leq \sqrt{\frac{3\Gamma(1-\alpha)e^{2|\lambda|T}T^\alpha(b-a)}{C_\Omega}} c_1(\tau^2 + h^4), \quad 1 \leq n \leq N$$

holds, where C_Ω is defined as that in Lemma 4.5.

Proof Subtracting (3.10)-(3.12) from (3.6), (3.8)-(3.9) respectively, we get the error equations

$$\begin{aligned}
\mathcal{A}_x \delta_t^{\alpha,\lambda} e_i^n - \mathcal{A}_t^\alpha \delta_x^2 e_i^n &= R_i^n, \quad 1 \leq i \leq M-1, \quad 1 \leq n \leq N, \\
e_0^n &= 0, \quad e_M^n = 0, \quad 1 \leq n \leq N, \\
e_i^0 &= 0, \quad 0 \leq i \leq M.
\end{aligned}$$

It follows from Theorem 4.6 and the truncation error estimate (3.7) that

$$\begin{aligned}
\|e^n\|^2 &\leq \frac{3\Gamma(1-\alpha)e^{2|\lambda|T}T^\alpha}{C_\Omega} \max_{0 \leq n \leq N} \|R^n\|^2 \\
&\leq \frac{3\Gamma(1-\alpha)e^{2|\lambda|T}T^\alpha(b-a)}{C_\Omega} c_1^2(\tau^2 + h^4)^2, \quad 1 \leq n \leq N.
\end{aligned}$$

This completes the proof. \square

5. An improved algorithm

This section presents an improved algorithm for the problem (1.2)-(1.4) with nonsmooth solution. To begin with this section, we first introduce the following lemma.

Lemma 5.1. *Let $f(t) = e^{-\lambda t} t^\beta$, $\beta > 0$. Then*

$$(1 - \frac{\alpha}{2})[-\infty D_t^{\alpha, \lambda} f(t)]_{t=t_n} + \frac{\alpha}{2}[-\infty D_t^{\alpha, \lambda} f(t)]_{t=t_{n-1}} = \frac{1}{\tau^\alpha} \sum_{k=0}^n g_k^{\alpha, \lambda} f(t_{n-k}) + \mathcal{O}(t_n^{\beta-\alpha-2} \tau^2)$$

*uniformly **holds** in $t \in \mathbb{R}$ as $\tau \rightarrow 0$, where $g_k^{\alpha, \lambda} = e^{-(k-\frac{\alpha}{2})\lambda\tau} g_k^\alpha$.*

The proof of above lemma is similar to [25] **and** here we skip it.

To fix the idea, let us first consider the fractional ordinary differential equation with substantial derivative as below

$${}_0 D_t^{\alpha, \lambda} u(t) = \mu u(t) + F(t). \quad (5.1)$$

We can seek its solution of the following form

$$u(t) = \sum_{j,k=0}^{+\infty} c_{j,k} e^{-\lambda t} t^{k+j\alpha}, \quad (5.2)$$

which can be derived by the method of power series. Indeed, suppose that the right-hand side function F is sufficiently smooth and has the power series expansion of the form below

$$F(t) = \sum_{k=0}^{+\infty} F_k e^{-\lambda t} t^k. \quad (5.3)$$

Inserting (5.2) and (5.3) into (5.1) leads to

$$\sum_{j,k=0}^{+\infty} c_{j,k} {}_0 D_t^{\alpha, \lambda} (e^{-\lambda t} t^{k+j\alpha}) = \mu \sum_{j,k=0}^{+\infty} c_{j,k} e^{-\lambda t} t^{k+j\alpha} + \sum_{k=0}^{+\infty} F_k e^{-\lambda t} t^k.$$

Consequently

$$\begin{aligned} & \sum_{j,k=0}^{+\infty} c_{j,k} e^{-\lambda t} \frac{\Gamma(k+1+j\alpha)}{\Gamma(k+1+(j-1)\alpha)} t^{k+(j-1)\alpha} \\ &= \mu \sum_{j,k=0}^{+\infty} c_{j,k} e^{-\lambda t} t^{k+j\alpha} + \sum_{k=0}^{+\infty} F_k e^{-\lambda t} t^k, \end{aligned} \quad (5.4)$$

where we have used the formula

$${}_0D_t^{\alpha,\lambda}(e^{-\lambda t}t^{k+j\alpha}) = e^{-\lambda t} \frac{\Gamma(k+1+j\alpha)}{\Gamma(k+1+(j-1)\alpha)} t^{k+(j-1)\alpha}.$$

Comparing the coefficients of both sides of identity (5.4) can solve $c_{j,k}$, the coefficients to be determined.

Now we turn our attention back to the TFSDE (1.2)-(1.4). From the aforementioned discussion, it makes sense to suppose the solution has the following structure:

$$u(\mathbf{x}, t) = \sum_{j,k=0}^{+\infty} c_{j,k}(\mathbf{x}) e^{-\lambda(\mathbf{x})t} t^{k+j\alpha}. \quad (5.5)$$

From the asymptotic behavior of the solution near the original point $t = 0$ in (5.5), we can clearly see, by Lemma 5.1, that the expected convergence order of the scheme proposed in the previous section may decrease correspondingly and thus the accuracy will deteriorate meantime. To make up for the **lost** accuracy, we have to turn to other approaches. One of the natural ideas occurs to us: if we can remove or separate the singular part from the candidate solution, then the remaining part has desired regularity to satisfy the necessary smoothness assumptions. However, this seems to be ideal but can not work in practice since the coefficients $c_{j,k}$ is unknown and difficult to calculate. An efficient approach firstly proposed by Lubich [25] is to add starting quadrature to make sure that the corrected scheme is accurate for the power function t^β , $\beta \in \Theta$, where $\Theta = \{\beta \mid \beta = k + j\alpha : j, k \in \mathbb{N}, \beta \leq 2 + \alpha\}$. Following the spirit of Lubich [25], by Lemma 5.1 we get the following semi-discrete correction scheme

$$\delta_t^{\alpha,\lambda} u(\mathbf{x}, t_n) + \sum_{k=0}^S w_{n,k}^{\alpha,\lambda} u(\mathbf{x}, t_k) = \mathcal{A}_t^\alpha \Delta u(\mathbf{x}, t_n) + \mathcal{A}_t^\alpha F(\mathbf{x}, t_n) + \mathcal{O}(\tau^2), \quad 1 \leq n \leq N, \quad (5.6)$$

for the problem (1.2)-(1.4) with solution in the form of (5.5), where S is the number of elements of the set Θ and the starting weights $w_{n,k}^{\alpha,\lambda}$ can be computed from the Vandermonde type system

$$\begin{aligned} \sum_{k=0}^S w_{n,k}^{\alpha,\lambda} t_k^{\beta_j} e^{-\lambda t_k} &= \left(1 - \frac{\alpha}{2}\right) \frac{\Gamma(\beta_j + 1)}{\Gamma(\beta_j + 1 - \alpha)} t_n^{\beta_j - \alpha} e^{-\lambda t_n} \\ &+ \frac{\alpha}{2} \frac{\Gamma(\beta_j + 1)}{\Gamma(\beta_j + 1 - \alpha)} t_{n-1}^{\beta_j - \alpha} e^{-\lambda t_{n-1}} - \frac{1}{\tau^\alpha} \sum_{k=0}^n g_k^{\alpha,\lambda} t_n^{\beta_j} e^{-\lambda t_k}, \quad j = 0, \dots, S. \end{aligned} \quad (5.7)$$

Instead of studying the theory of the improved algorithm, we will provide a two-dimensional example to show its validity in next section, where a fully

discretized compact finite scheme for the two-dimensional TFSDE is also given; see Example 6.3.

Remark 5.2. *Based on the pitfalls in fast numerical solvers for fractional differential equations studied in [15], it is known that the linear system (5.7) is ill-posed, which may result in the reduced efficiency of the algorithm (5.6). However, using a few correction terms can significantly increase the accuracy for problems with low regularity. It will not be computational expensive to calculate the necessary starting weights; see also [43].*

6. Numerical examples

In this section, we give some numerical results to illustrate the performance of the proposed scheme and verify the theoretical prediction. In Example 6.1, a prototypical equation is considered to show the convergence order of the proposed second-order approximation (2.9) with or without the smoothness assumptions in Theorem 2.3. Example 6.2 provides a comparison between the proposed scheme (3.10)-(3.12) and the scheme in [14], for solving the backward Feynman-Fac equation (6.1). In Example 6.3, we illustrate efficiency of the improved algorithm by solving a two-dimensional TFSDE with nonsmooth solution.

Example 6.1. *Consider the following prototypical equation*

$${}_0D_t^{\alpha,\lambda}u(t) = f(t),$$

with $u(0)$ and the source term $f(t)$ such that the equation admits the solution $u(t) = e^{-\lambda t}(t^3 + t^\nu)$, and $0 < \alpha < 1$. We take $\lambda = 0.5$ and change ν to test the convergence order of the approximation (2.9) for the problem with different regularity.

Denote

$$E(\tau) = \max_{1 \leq n \leq N} |u^n - U^n|.$$

In this example $\text{Rate} := \log_2 \frac{E(\tau)}{E(\tau/2)}$.

Tables 6.1 displays that the approximation (2.9) can obtain second-order accuracy in uniform maximum norm when $\nu = 2.5$ and 2, while the convergence

Table 6.1: Error and convergence order of the approximation (2.9) for $u(t)$ with different regularity (Example 6.1).

| | τ | $\alpha = 0.2$ | | $\alpha = 0.5$ | | $\alpha = 0.8$ | |
|-------------|--------|----------------|------|----------------|------|----------------|------|
| | | $E(\tau)$ | Rate | $E(\tau)$ | Rate | $E(\tau)$ | Rate |
| $\nu = 2.5$ | 1/16 | 1.9225e-4 | | 4.7884e-4 | | 7.5901e-4 | |
| | 1/32 | 4.8101e-5 | 2.00 | 1.2002e-4 | 1.99 | 1.9083e-4 | 2.00 |
| | 1/64 | 1.2029e-5 | 2.00 | 3.0043e-5 | 2.00 | 4.7871e-5 | 2.00 |
| | 1/128 | 3.0076e-6 | 2.00 | 7.5152e-6 | 2.00 | 1.1993e-5 | 2.00 |
| $\nu = 2$ | 1/16 | 1.5725e-4 | | 3.8400e-4 | | 5.6264e-4 | |
| | 1/32 | 3.9392e-5 | 2.00 | 9.6827e-5 | 1.99 | 1.4297e-4 | 1.98 |
| | 1/64 | 9.8586e-6 | 2.00 | 2.4348e-5 | 1.99 | 3.6235e-5 | 1.98 |
| | 1/128 | 2.4661e-6 | 2.00 | 6.1116e-6 | 1.99 | 9.1649e-6 | 1.98 |
| $\nu = 1.5$ | 1/16 | 1.3578e-4 | | 3.1355e-4 | | 3.0475e-4 | |
| | 1/32 | 3.6556e-5 | 1.89 | 7.8536e-5 | 2.00 | 1.2018e-4 | 1.34 |
| | 1/64 | 1.2691e-5 | 1.53 | 1.9651e-5 | 2.00 | 4.4176e-5 | 1.44 |
| | 1/128 | 4.4625e-6 | 1.51 | 4.9148e-6 | 2.00 | 1.5849e-5 | 1.48 |
| $\nu = 1$ | 1/16 | 8.0608e-4 | | 3.1355e-4 | | 1.0800e-3 | |
| | 1/32 | 4.0503e-4 | 0.99 | 7.0492e-4 | 1.00 | 5.2210e-4 | 1.05 |
| | 1/64 | 2.0355e-4 | 0.99 | 3.5386e-4 | 0.99 | 2.6071e-4 | 1.00 |
| | 1/128 | 1.0211e-4 | 1.00 | 1.7745e-4 | 1.00 | 1.3082e-4 | 0.99 |
| $\nu = 0.5$ | 1/16 | 1.0492e-2 | | 2.7597e-2 | | 4.9525e-2 | |
| | 1/32 | 7.5289e-3 | 0.48 | 1.9804e-2 | 0.48 | 3.5543e-2 | 0.48 |
| | 1/64 | 5.3646e-3 | 0.49 | 1.4111e-2 | 0.49 | 2.5327e-2 | 0.49 |
| | 1/128 | 3.8081e-3 | 0.49 | 1.0017e-2 | 0.49 | 1.7978e-2 | 0.49 |

order decays with the decrease of ν evidently. The reduction of accuracy implies that some additional assumptions are needed to get second-order convergence for the approximation (2.9), e.g., the vanishing derivatives $u_t(0) = 0$. To keep the second-order convergence of the approximation (2.9), we make an assumption of the extended solution $\tilde{u}(t) \in \mathcal{C}^{2+\alpha}(\mathbb{R})$ in this paper. To some extent the assumption may be too strict, so we have considered the use of correction terms in the proposed schemes for problems with low regularity; see Section 5 for the improved schemes and Example 6.3 for the corresponding numerical results.

Example 6.2. Consider the backward fractional Feynman-Kac equation presented in [14]

$${}_0\partial_t^{\alpha, \lambda(x)}[P(x, \rho, t) - e^{-\rho U(x)t}P(x, \rho, 0)] = \kappa_\alpha \partial_x^2 P(x, \rho, t) + f(x, t) \quad (6.1)$$

on a finite domain $0 < x < 1$, $0 < t \leq 1$, with the coefficients $\kappa_\alpha = 0.5$, and

$\lambda(x) = \rho U(x)$, $U(x) = x$, $\rho = 1 + \sqrt{-1}$; the forcing function

$$f(x, t) = -\kappa_\alpha e^{-\rho x t} (t^{3+\alpha} + 1) (\rho^2 t^2 \sin(\pi x) - 2\pi \rho t \cos(\pi x) - \pi^2 \sin(\pi x)), \\ + \frac{\Gamma(4 + \alpha)}{\Gamma(4)} e^{-\rho x t} t^3 \sin(\pi x). \quad (6.2)$$

the initial condition $P(x, \rho, 0) = \sin(\pi x)$, and the boundary conditions $P(0, \rho, t) = P(1, \rho, t) = 0$. The exact solution is given by $P(x, \rho, t) = e^{-\rho x t} (t^{3+\alpha} + 1) \sin(\pi x)$.

Notice that the initial value is not equal to zero. So we have made the transformation

$$u(x, t) = P(x, \rho, t) - e^{-\rho x t} P(x, \rho, 0).$$

Table 6.2: Error and convergence order of the scheme (3.10)-(3.12) in time with a fixed space step-size $h = 1/40$ and $\lambda(x) = (1 + \sqrt{-1})x$ (Example 6.2).

| τ | $\alpha = 0.2$ | | $\alpha = 0.5$ | | $\alpha = 0.8$ | |
|--------|----------------|------|----------------|------|----------------|------|
| | $E_1(\tau, h)$ | Rate | $E_1(\tau, h)$ | Rate | $E_1(\tau, h)$ | Rate |
| 1/5 | 5.1150e-3 | | 1.3602e-2 | | 2.1793e-2 | |
| 1/10 | 1.3025e-3 | 1.97 | 3.4574e-3 | 1.98 | 5.5020e-3 | 1.99 |
| 1/20 | 3.2845e-4 | 1.99 | 8.7149e-4 | 1.99 | 1.3819e-3 | 1.99 |
| 1/40 | 8.2375e-5 | 2.00 | 2.1870e-4 | 1.99 | 3.4619e-4 | 2.00 |

Table 6.3: Error and convergence order of the scheme (3.10)-(3.12) in space with a fixed time step $\tau = 1/10000$ and $\lambda(x) = (1 + \sqrt{-1})x$ (Example 6.2).

| h | $\alpha = 0.2$ | | $\alpha = 0.5$ | | $\alpha = 0.8$ | |
|------|----------------|------|----------------|------|----------------|------|
| | $E_1(\tau, h)$ | Rate | $E_1(\tau, h)$ | Rate | $E_1(\tau, h)$ | Rate |
| 1/4 | 2.0681e-03 | | 1.7956e-03 | | 1.4439e-03 | |
| 1/8 | 1.2566e-04 | 4.04 | 1.1059e-04 | 4.02 | 9.0999e-05 | 3.99 |
| 1/16 | 7.9126e-06 | 3.99 | 6.9117e-06 | 4.00 | 5.6411e-06 | 4.01 |
| 1/32 | 4.9271e-07 | 4.01 | 4.2904e-07 | 4.01 | 3.4901e-07 | 4.01 |
| 1/64 | 2.9927e-08 | 4.04 | 2.4756e-08 | 4.12 | 1.8798e-08 | 4.21 |

Actually, replacing the average operators \mathcal{A}_t^α and \mathcal{A}_x in (3.10)-(3.11) by identity operators in temporal and spatial directions respectively, we can obtain the following scheme in [14] as below

$$\delta_t^{\alpha, \lambda_i} u_i^n - \delta_x^2 u_i^n = F_i^n, \quad 1 \leq i \leq M-1, \quad 1 \leq n \leq N, \quad (6.3)$$

$$u_i^0 = u_0(x_i), \quad 0 \leq i \leq M, \quad (6.4)$$

$$u_0^n = \phi_a(x_0, \mathbf{t}_n), \quad u_M^n = \phi_b(x_M, \mathbf{t}_n), \quad 1 \leq n \leq N. \quad (6.5)$$




Figure 6.1: Convergence order and error comparison between the scheme (3.10)-(3.12) and the scheme (6.3)-(6.5) in [14] with $\alpha = 0.5$ for solving Equation (6.1) at $t = 1$. In all cases, we set $N = M^2$ (Example 6.2).

Define the error

$$E_1(\tau, h) = \max_{1 \leq i \leq M-1} |u(x_i, t_N) - u_i^N|,$$

where $u(x_i, t_N)$ represents the exact solution and u_i^N is the numerical solution with the mesh step-sizes h and τ at the grid point (x_i, t_N) . Assume

$$E_1(\tau, h) = O(\tau^p) + O(h^q).$$

If τ is sufficiently small, then $E_1(\tau, h) \approx O(h^q)$. Consequently, $\frac{E_1(\tau, 2h)}{E_1(\tau, h)} \approx 2^q$ and $q \approx \log_2 \left(\frac{E_1(\tau, 2h)}{E_1(\tau, h)} \right)$ is the convergence order with respect to the spatial step size. Likewise, the convergence order in time can be taken as $p \approx \log_2 \left(\frac{E_1(\tau, 2h)}{E_1(\tau, h)} \right)$ with sufficiently small h . Tables 6.2-6.3 list the errors and convergence orders of our scheme, showing that it is of second-order convergence in time and fourth-order convergence in space respectively. This is in a good agreement with our theoretical results. Evidently, we can see from Figure 6.1 that our proposed scheme displays much higher accuracy and enjoys advantage over the scheme (6.3)-(6.4) presented in [14]. Table 6.4 further shows that when the two schemes generate the same accuracy, our proposed scheme needs fewer temporal and spatial grid points and less CPU time than that of scheme (6.3)-(6.4).

Example 6.3. Consider the following two dimensional backward fractional

Table 6.4: A comparison between the scheme (3.10)-(3.12) and the scheme (6.3)-(6.5) in [14] for different α and step sizes for solving Equation (6.1) (Example 6.2).

| | the scheme (3.10)-(3.12) | | | | the scheme (6.3)-(6.5) | | | |
|----------------|--------------------------|-----|----------------|-------------|------------------------|-----|----------------|-------------|
| | N | M | $E_1(\tau, h)$ | CPU(second) | N | M | $E_1(\tau, h)$ | CPU(second) |
| $\alpha = 0.1$ | 16 | 4 | 1.9804e-3 | 0.0151 | 256 | 16 | 2.2831e-3 | 1.0427 |
| | 36 | 6 | 3.8109e-4 | 0.0340 | 1296 | 36 | 4.5106e-4 | 50.9968 |
| | 64 | 8 | 1.1946e-4 | 0.0661 | 4096 | 64 | 1.4269e-4 | 1484.3471 |
| $\alpha = 0.5$ | 16 | 4 | 1.2906e-3 | 0.0090 | 256 | 16 | 2.3822e-3 | 0.9892 |
| | 36 | 6 | 2.5103e-4 | 0.0262 | 1296 | 36 | 4.6994e-4 | 50.6735 |
| | 64 | 8 | 7.9988e-5 | 0.0671 | 4096 | 64 | 1.4863e-4 | 1491.8938 |
| $\alpha = 0.9$ | 16 | 4 | 1.6037e-3 | 0.0077 | 256 | 16 | 2.8498e-3 | 0.9932 |
| | 36 | 6 | 3.4057e-4 | 0.0261 | 1296 | 36 | 5.6214e-4 | 51.8453 |
| | 64 | 8 | 1.0771e-4 | 0.0687 | 4096 | 64 | 1.7779e-4 | 1471.3060 |

Table 6.5: Error and convergence order of the scheme (6.7)-(6.9) for the 2D problem (6.6). $\alpha = 0.2$, $h_1 = h_2 = 1/60$; S denotes the number of correction terms (Example 6.3).

| N | $S = 0$ | | $S = 1$ | | $S = 2$ | | $S = 3$ | |
|-----|-----------------------|------|-----------------------|------|-----------------------|------|-----------------------|------|
| | $E_2(\tau, h_1, h_2)$ | Rate | $E_2(\tau, h_1, h_2)$ | Rate | $E_2(\tau, h_1, h_2)$ | Rate | $E_2(\tau, h_1, h_2)$ | Rate |
| 4 | 9.32e-3 | | 8.86e-4 | | 6.68e-4 | | 1.47e-3 | |
| 8 | 8.68e-3 | 0.10 | 2.76e-4 | 1.68 | 1.95e-4 | 1.78 | 2.34e-4 | 2.65 |
| 16 | 8.03e-3 | 0.11 | 2.02e-4 | 0.45 | 5.09e-5 | 1.94 | 5.31e-5 | 2.14 |
| 32 | 7.43e-3 | 0.11 | 1.70e-4 | 0.25 | 1.32e-5 | 1.95 | 1.33e-5 | 2.00 |

Feynman-Kac equations

$${}_0\partial_t^{\alpha, \lambda(x, y)}[u(x, y, t) - e^{-\lambda(x, y)t}u(x, y, 0)] = \Delta u(x, y, t) + F(x, y, t) \quad (6.6)$$

on a finite domain $\Omega = (0, 1) \times (0, 1)$, $0 < t \leq 1$, and $\lambda(x, y) = 0.01(x + y)$. We take the initial condition $u(x, y, 0) = \sin(\pi x) \sin(\pi y)$, and the boundary condition $u(x, y, t) = 0$ for $(x, y) \in \partial\Omega$. The exact solution is

$$u(x, y, t) = e^{-\lambda(x, y)t}(1 + t^\alpha + t^{2\alpha} + t^3) \sin(\pi x) \sin(\pi y).$$

For spatial approximation, take two integers M_1, M_2 and let $h_1 = 1/M_1$, $h_2 = 1/M_2$, $x_i = ih_1$, $0 \leq i \leq M_1$, $y_j = jh_2$, $0 \leq j \leq M_2$. Let $\bar{\Omega}_h = \{(x_i, y_j) | 0 \leq i \leq M_1, 0 \leq j \leq M_2\}$, and $\Omega_h = \bar{\Omega}_h \cap \Omega$, and $\partial\Omega_h = \bar{\Omega}_h \cap \partial\Omega$. For any grid function $v = \{v_{i,j} | 0 \leq i \leq M_1, 0 \leq j \leq M_2\}$, denote

$$\delta_x v_{i-\frac{1}{2}, j} = \frac{1}{h_1}(v_{i,j} - v_{i-1,j}), \quad \delta_x^2 v_{i,j} = \frac{1}{h_1}(\delta_x v_{i+\frac{1}{2}, j} - \delta_x v_{i-\frac{1}{2}, j}).$$

Similar notations $\delta_y v_{i,j-\frac{1}{2}}$, $\delta_y^2 v_{i,j}$ can be defined. The spatial average operators are defined as

$$\mathcal{A}_x v_{i,j} = \begin{cases} \frac{1}{12}(v_{i-1,j} + 10v_{i,j} + v_{i+1,j}), & 1 \leq i \leq M_1 - 1, \quad 0 \leq j \leq M_2, \\ v_{i,j}, & i = 0 \text{ or } M_1, \quad 0 \leq j \leq M_2, \end{cases}$$

$$\mathcal{A}_y v_{i,j} = \begin{cases} \frac{1}{12}(v_{i,j-1} + 10v_{i,j} + v_{i,j+1}), & 1 \leq j \leq M_2 - 1, \quad 0 \leq i \leq M_1, \\ v_{i,j}, & j = 0 \text{ or } M_2, \quad 0 \leq i \leq M_1. \end{cases}$$

By generalizing the improved scheme (5.6), we present a compact difference scheme with correction terms in time for the two-dimensional equation (6.6)

$$\mathcal{A}_x \mathcal{A}_y [\delta_t^{\alpha,\lambda} u_{i,j}^n + \sum_{k=0}^S w_{n,k}^{\alpha,\lambda} u_{i,j}^n] - \mathcal{A}_t^\alpha \mathcal{A}_y \delta_x^2 u_{i,j}^n - \mathcal{A}_t^\alpha \mathcal{A}_x \delta_y^2 u_{i,j}^n = \mathcal{A}_t^\alpha \mathcal{A}_x \mathcal{A}_y F_{i,j}^n,$$

$$(x_i, y_j) \in \Omega_h, \quad 1 \leq n \leq N, \quad (6.7)$$

$$u_{i,j}^0 = u_0(x_i, y_j), \quad (x_i, y_j) \in \bar{\Omega}_h, \quad (6.8)$$

$$u_{i,j}^n = \phi(x_i, y_j, \textcolor{red}{t}_n), \quad (x_i, y_j) \in \partial\Omega_h, \quad 1 \leq n \leq N. \quad (6.9)$$

Denote

$$E_2(\tau, h_1, h_2) = \max_{1 \leq n \leq N} \|u(\cdot, \cdot, \textcolor{red}{t}_n) - u^n\|_\infty.$$

Table 6.5 shows that the errors behave well and convergence order is augmented up to two with the increase of the number of correction terms, which indicates the efficiency and validity of our improved algorithm.

7. Conclusion

We have proposed a second-order approximation for the fractional substantial derivative and applied it to solve the time-fractional substantial diffusion equation. By combining the fourth-order compact finite difference approximation, a fully discrete Grünwald-Letnikov-formula-based compact difference scheme has been presented. It has been proved that the proposed scheme is unconditionally stable and of second-order convergence in time and fourth-order convergence in space.

We have numerically showed that our proposed scheme can reach the predicted accuracy when solving a problem with smooth solution. Compared with the scheme proposed in [14] for solving the backward Feymann-Fac fractional equation, our scheme requires less storage and computational cost for the same accuracy. While if the solution is not smooth enough, we have illustrated in Example 6.1 that our proposed scheme loses its accuracy to some extent. To overcome this difficulty, we followed the idea of Lubich [25] and introduced the

corresponding starting quadratures of substantial version. We have proposed an improved algorithm by applying the correction terms to increase the accuracy of the scheme near the origin and recover the second-order accuracy in time for problems with nonsmooth solution, which has been demonstrated by Example 6.3.

By the modified Grünwald derivative and the shifted Grünwald-Letnikov-formula, we have presented a weighted and shifted substantial Grünwald formula and given its asymptotic expansion, which is essential to derive the second-order approximation for the fractional substantial derivative. The proposed weighted average method is an efficient tool to get high-order scheme, not limited to the problem under consideration in this study. **It can also be extended and applied to solve tempered fractional equations, which have attracted considerable interest recently; see [12, 40].** It should be also pointed out that our scheme is consistent with the Crank-Nicolson scheme when the fractional derivatives reduces to the standard operators, which is one of attractive features of our proposed scheme.

Acknowledgement

The research is partially supported by National Natural Science Foundation of China (No. 11271068) and by the Fundamental Research Funds for the Central Universities and the Research and Innovation Project for College Graduates of Jiangsu Province (Grant No.: KYLX_0081). G. Lin would like to thank the support by NSF Grant DMS-1115887, and the U.S. Department of Energy, Office of Science, Office of Advanced Scientific Computing Research, Applied Mathematics program as part of the Collaboratory on Mathematics for Mesoscopic Modeling of Materials, and Multifaceted Mathematics for Complex Energy Systems.

References

- [1] A. A. Alikhanov, A new difference scheme for the time fractional diffusion equation, *J. Comput. Phys.*, 280 (2015) 424–438.
- [2] B. Baeumer, M. M. Meerschaert, Tempered stable Lévy motion and transient super-diffusion, *J. Comput. Appl. Math.*, 233 (2010) 2438–2448.
- [3] I. Bronstein, Y. Israel, E. Kepten, S. Mai, Y. Shav-Tal, E. Barkai, Y. Garini, Transient anomalous diffusion of telomeres in the nucleus of mammalian cells, *Phys. Rev. Lett.*, 103 (2009) 018102.

- [4] R. G. Buschman, Decomposition of an integral operator by use of Mikusiński calculus, *SIAM J. Math. Anal.*, 3 (1972) 83–85.
- [5] G. G. Cabal, A. Genovesio, S. R.-Navarro, C. Zimmer, O. Gadai, A. Lesne, H. Buc, F. F.-Fournier, J.-C. O.-Marin, E. C. Hurt, U. Nehrbass, SAGA interacting factors confine sub-diffusion of transcribed genes to the nuclear envelope, *Nature*, 441 (2006) 770–773.
- [6] J. Cao , C. Li , Y.Q. Chen , On tempered and substantial fractional calculus, *Mechatronic and Embedded Systems and Applications (MESA)*, 2014 IEEE/ASME 10th International Conference on., IEEE, (2014) 1-6.
- [7] W. Cao, Z. Zhang, G. E. Karniadakis, Time-splitting schemes for fractional differential equations I: smooth solutions, *SIAM J. Sci. Comput.*, 37(4) (2015) A1752-A1776.
- [8] S. Carmi, E. Barkai, Fractional Feynman-Kac equation for weak ergodicity breaking, *Phys. Rev. E*, 84 (2011) 061104.
- [9] S. Carmi, L. Turgeman, E. Barkai, On distributions of functionals of anomalous diffusion paths, *J. Statis. Phys.*, 141 (2010) 1071–1092.
- [10] A. Cartea, D. del Castillo-Negrete, Fluid limit of the continuous-time random walk with general Levy jump distribution functions, *Phys. Rev. E*, 76 (2007) 041105.
- [11] M. Chen, W. Deng, Discretized fractional substantial calculus, *ESAIM-Math. Model. Num.*, 49 (2015) 373-394.
- [12] M. Chen, W. Deng, High order algorithms for the fractional substantial diffusion equation with truncated Lévy flights, *SIAM J. Sci. Comput.*, 37 (2015) A890–A917.
- [13] A. Comtet, J. Desbois, C. Texier, Functionals of Brownian motion, localization and metric graphs, *J. Phys. A-Math. General*, 38 (2005) R341–R383.
- [14] W. Deng, M. Chen, E. Barkai, Numerical algorithms for the forward and backward fractional Feynman–Kac equations, *J. Sci. Comput.*, 62 (2015) 718–746.
- [15] K. Diethelm, N. J. Ford, A. D. Freed, and M. Weilbeer, Pitfalls in fast numerical solvers for fractional differential equations, *J. Comput. Appl. Math.*, 186 (2006) 482-503.

- [16] Y. Dimitrov, Numerical approximations for fractional differential equations, *J. Fract. Calc. Appl.*, 5 (2014) 45.
- [17] R. Friedrich, F. Jenko, A. Baule, S. Eule, Anomalous diffusion of inertial, weakly damped particles, *Phys. Rev. Lett.*, 96 (2006) 230601.
- [18] G. Gao, H. Sun, Z. Sun, Stability and convergence of finite difference schemes for a class of time-fractional sub-diffusion equations based on certain superconvergence, *J. Comput. Phys.*, 280 (2015) 510–528.
- [19] E. Hanert, C. Piret, A Chebyshev pseudospectral method to solve the space-time tempered fractional diffusion equation, *SIAM J. Sci. Comput.*, 36 (2014) A1797–A1812.
- [20] B. Jin, R. Lazarov, Z. Zhou, Error estimates for a semidiscrete finite element method for fractional order parabolic equations, *SIAM J. Numer. Anal.*, 51 (2013) 445–466.
- [21] D. Kleinhans, R. Friedrich, Continuous-time random walks: Simulation of continuous trajectories, *Phys. Rev. E*, 76 (2007) 061102.
- [22] A. Kullberg, D. del C.-Negrete, Transport in the spatially tempered, fractional Fokker-Planck equation, *J. Phys. A-Math. Theor.*, 45 (2012) 255101.
- [23] Y. Lin, C. Xu, Finite difference/spectral approximations for the time-fractional diffusion equation, *J. Comput. Phys.*, 225 (2007) 1533–1552.
- [24] F. Liu, P. Zhuang, I. Turner, K. Burrage, V. Anh, A new fractional finite volume method for solving the fractional diffusion equation, *Appl. Math. Model.*, 38 (2014) 3871–3878.
- [25] C. Lubich, Discretized fractional calculus, *SIAM J. Math. Anal.*, 17 (1986) 704–719.
- [26] M. M. Meerschaert, C. Tadjeran, Finite difference approximations for fractional advection-dispersion flow equations, *J. Comput. Appl. Math.*, 172 (2004) 65–77.
- [27] M. M. Meerschaert, Y. Zhang, B. Baeumer, Tempered anomalous diffusion in heterogeneous systems. *Geophys. Res. Lett.*, 35 (2008) 190201.
- [28] R. Metzler, J. Klafter, The random walk’s guide to anomalous diffusion: a fractional dynamics approach, *Phys. Rep.-Review Section of Physics Letters*, 339 (2000) 1–77.

- [29] K. B. Oldham, J. Spanier, *The fractional calculus*, Academic Press, New York-London, 1974.
- [30] I. Podlubny, *Fractional differential equations*, Academic Press, New York, 1999.
- [31] I. Podlubny, T. Skovranek, B. M. V. Jara, I. Petras, V. Verbitsky, Y. Chen, Matrix approach to discrete fractional calculus III: non-equidistant grids, variable step length and distributed orders, *Philos. Trans. R. Soc. Lond. Ser. A Math. Phys. Eng. Sci.*, 371 (2013) 20120153.
- [32] S. Shen, F. Liu, V. Anh, Numerical approximations and solution techniques for the space-time Riesz-Caputo fractional advection-diffusion equation, *Numer. Algorithms*, 56 (2011) 383–403.
- [33] I. Sokolov, R. Metzler, Towards deterministic equations for Levy walks: The fractional material derivative, *Phys. Rev. E*, (2003) 010101.
- [34] Z.Z. Sun, X. Wu, A fully discrete difference scheme for a diffusion-wave system, *Appl. Numer. Math.*, 56 (2006) 193–209.
- [35] Z.Z. Sun, *The method of order reduction and its application to the numerical solutions of partial differential equations*, Science Press, Beijing, 2009.
- [36] Z. Z. Sun, Numerical methods of partial differential equations, 2D Edition, Science Press, Beijing, 2012 (in Chinese).
- [37] C. Tadjeran, M. M. Meerschaert, H.-P. Scheffler, A second-order accurate numerical approximation for the fractional diffusion equation, *J. Comput. Phys.*, 213 (2006) 205–213.
- [38] L. Turgeman, S. Carmi, E. Barkai, Fractional Feynman-Kac equation for non-Brownian functionals. *Phys. Rev. Lett.*, 103 (2009) 190201.
- [39] S. B. Yuste, J. Q.-Murillo, A finite difference method with non-uniform timesteps for fractional diffusion equations, *Comput. Phys. Commun.*, 183 (2012) 2594–2600.
- [40] M. Zayernouri, M. Ainsworth, G. Karniadakis, Tempered fractional Sturm-Liouville eigenproblems, *SIAM J. Sci. Comput.* 37 (4) (2015) A1777-A1800.
- [41] F. Zeng, C. Li, F. Liu, I. Turner, The use of finite difference/element approaches for solving the time-fractional subdiffusion equation, *SIAM J. Sci. Comput.*, 35 (2013) A2976–A3000.

- [42] F. Zeng, C. Li, F. Liu, I. Turner, Numerical algorithms for time-fractional subdiffusion equation with second-order accuracy, *SIAM J. Sci. Comput.*, 37 (2015) A55–A78.
- [43] F. Zeng, Z. Zhang, G. Karniadakis, Second-order numerical methods for multi-term fractional differential equations: smooth and non-smooth solutions, *To appear*.